



West Nile Virus Disease Cases* and Presumptive Viremic Blood Donors by State – United States, 2020

State	Neuroinvasive Disease Cases†	Non-neuroinvasive	Total cases	Deaths	Presumptive Viremic Blood Donors‡
		Disease Cases			
Alabama	7	2	9	1	3
Alaska	0	0	0	0	0
Arizona	8	4	12	2	1
Arkansas	1	0	1	0	0
California	179	56	235	11	32
Colorado	17	18	35	1	8
Connecticut	5	3	8	0	0
Delaware	0	0	0	0	0
District of Columbia	0	0	0	0	0
Florida	44	7	51	3	41
Georgia	7	1	8	1	4
Hawaii	0	0	0	0	0
Idaho	5	1	6	0	1
Illinois	36	6	42	4	0
Indiana	3	0	3	0	0
lowa	2	1	3	0	0
Kansas	6	2	8	0	0
Kentucky	0	0	0	0	0
Louisiana	14	4	18	3	5
Maine	1	0	1	0	0
Maryland	<u>+</u> 1	0	1	0	0
Massachusetts	9	3	12	<u>0</u> 1	4
					3
Michigan	29	3	32	1	
Minnesota Mississippi	0	0	0	0	0
Mississippi	4	3	7	0	0
Missouri	1	0	1	0	0
Montana	1	0	1	0	0
Nebraska	9	6	15	1	3
Nevada	0	1	1	0	0
New Hampshire	0	0	0	0	0
New Jersey	3	0	3	0	1
New Mexico	7	1	8	1	0
New York	18	4	22	4	3
North Carolina	1	0	1	1	0
North Dakota	2	5	7	0	1
Ohio	3	0	3	0	0
Oklahoma	6	1	7	4	1
Oregon	0	1	1	0	0
Pennsylvania	7	4	11	1	0
Rhode Island	0	0	0	0	0
South Carolina	4	0	4	0	2
South Dakota	10	10	20	0	3
Tennessee	0	0	0	0	0
Texas	101	21	122	26	30
Utah	1	1	2	0	0
Vermont	0	0	0	0	0
Virginia	1	1	2	0	1
Washington	0	2	2	0	0
West Virginia	0	0	0	0	0
Wisconsin	5	0	5	0	0
Wyoming	1	0	1	0	0
Totals	559	172	731	66	147

^{*}Includes confirmed and probable cases

[†]E.g., meningitis, encephalitis, or acute flaccid paralysis





‡Presumptive viremic blood donors (PVDs) are people who had no symptoms at the time of donating blood through a blood collection agency, but whose blood tested positive when screened for the presence of West Nile virus. Some PVDs develop symptoms after donation; of the 147 PVDs, 20 (14%) developed clinical illness and are also included as disease cases.